

Putting together and taking apart: assembly and disassembly of the Rad51 nucleoprotein filament in DNA repair and genome stability

Tadas Andriuskevicius¹, Oleksii Kotenko¹ and Svetlana Makovets^{1,*}

¹ Institute of Cell Biology, School of Biological Sciences, University of Edinburgh.

* Corresponding Author:

Svetlana Makovets, Institute of Cell Biology, School of Biological Sciences, University of Edinburgh, Alexander Crum Brown Road, King's Buildings, Edinburgh, EH9 3FF, United Kingdom; E-mail: Sveta.Makovets@ed.ac.uk

ABSTRACT Homologous recombination is a key mechanism providing both genome stability and genetic diversity in all living organisms. Recombinases play a central role in this pathway: multiple protein subunits of Rad51 or its orthologues bind single-stranded DNA to form a nucleoprotein filament which is essential for initiating recombination events. Multiple factors are involved in the regulation of this step, both positively and negatively. In this review, we discuss Rad51 nucleoprotein assembly and disassembly, how it is regulated and what functional significance it has in genome maintenance.

doi: 10.15698/cst2018.05.134

Received originally: 01.02.2018

in revised form: 12.03.2018,

Accepted 13.03.2018,

Published 28.03.2018.

Keywords: homologous recombination, Rad51 filament, double-stranded DNA break, DNA repair, Rad51 regulation.

Abbreviations:

BIR – break-induced replication,
DNMT – de novo telomere addition,
DSB – double-stranded DNA break,
dsDNA – double-stranded DNA,
GC – gene conversion,
HR – homologous recombination,
PTM – post-translational modification,
RFC – Replication Factor C,
RPA – Replication Protein A,
SDSA – synthesis-dependent strand annealing,
SIM – SUMO-interacting motif,
SSA – single-stranded annealing,
ssDNA – single-stranded DNA.

INTRODUCTION

Homologous recombination (HR) involves exchange of genetic information, often between two different DNA molecules. This exchange requires physical interaction between the molecules and may lead to heritable genetic changes, contributing to biodiversity and higher evolutionary adaptability of species. Novel combinations of alleles arising in meiosis is a great example of genetic variability as a result of HR. On the other hand, physical linkage between chromosomes during meiotic HR is important for pairing and accurate segregation of homologous chromosomes in meiosis, thereby preventing aneuploidy and promoting genome integrity. In addition, HR machinery contributes to

genome stability by playing a central role in stabilisation and restart of stalled replication forks as well as repair of DNA breaks routinely arising from broken replication forks, oxidative damage, etc. Because DNA breaks are part of normal cell physiology, unsurprisingly, defects in HR in humans lead to developmental disorders and cancer predisposition [1].

A double-stranded DNA break (DSB) is often considered the most dangerous type of DNA damage as it disrupts the continuity of chromosomes and, if unrepaired, may lead to loss of genetic information and eventual cell death. Two major mechanisms are normally used to repair a DSB: non-homologous end joining (NHEJ) and HR (**Figure 1**). NHEJ

involves ligation of the broken ends with little or no DNA processing around the break. It is efficient but often mutagenic due to small deletions or insertions at the damage locus [2-4]. Occasionally, it can also lead to gross chromosomal rearrangements due to illegitimate joining of DNA ends from different breaks or ligations of chromosome breaks to telomeres [5-7]. HR includes a group of pathways sharing two features: i) they require intact homologous DNA sequences for the repair (called donor DNA) and ii) they stem from the same original step of extensive DSB resection around the break, which generates single-stranded DNA (ssDNA) necessary for HR pathways to oper-

ate. Although DSB repair is well-conserved among eukaryotes, the preference for the repair mechanism is cell cycle-dependent and species-specific [3, 8, 9]. Occasionally, a DSB can be mistakenly recognised as a telomere and telomerase, the enzyme responsible for telomere extension, heals the break by adding a *de novo* telomere (**Figure 1**) [10]. Although *de novo* telomere addition (DNTE) may stabilise the broken chromosome, it results in terminal deletions.

Depending on the nature of a DSB break, homology-dependent repair may proceed through gene conversion (GC), break-induced replication (BIR) or single-strand an-




FIGURE 1: A general overview of DSB repair mechanisms. A DNA molecule with a DSB (blue lines) can either be repaired by NHEJ (A) or resected, thereby committing to HR. Sometimes, resected breaks can be healed by telomerase (DNTE, B) rather than repaired by HR but these events are rare. Resected DNA can be repaired by SSA (C) when there are direct repeats flanking the break (pink lines). Alternatively, resected ssDNA might invade a homologous donor molecule (red lines) enabling repair by BIR (D) or GC (E). During BIR, the invading strand is extended and the newly synthesised DNA is displaced from the donor to act as a template for the second strand. GC can proceed via SDSA or DSBR pathways. During SDSA, the invading ssDNA is extended, displaced from the donor and annealed to the other end of the break. DSBR involves the extension of the invading strand, capture of the second end of the break and the resolution of the dHJ intermediate formed. Arrow heads indicate 3' ends of the DNA. Dotted lines show newly-synthesised DNA and their colours correspond to the DNA molecules that have been used as templates. The black zigzag represents the telomere. Abbreviations: NHEJ – non-homologous end joining; DNTE – *de novo* telomere addition; SSA – single-strand annealing; BIR – break-induced replication; GC – gene conversion; SDSA – synthesis-dependent strand annealing; DSBR – double-strand break repair.

nealing (SSA; **Figure 1**). GC is used to repair a DSB when both ends of the break have homology to a donor DNA molecule and are available for the repair, whereas BIR is employed when only one end is present, for example, to restore broken replication forks [11, 12]. SSA does not involve external homology and can only be used if a DNA break is flanked by homologous repeats in direct orientation. SSA is mutagenic as any sequences between the homologies and one of the two copies of the repeated sequences are lost [13]. However, SSA may act as a safeguard mechanism when other options are not available.

All three homology-dependent repair pathways begin with the resection of the 5' end around a DSB to create ssDNA overhangs [14]. SSA may occur at this point through annealing of the complementary sequences on the resected ends [13]. GC and BIR require invasion of the resected DNA end into a donor molecule and a subsequent strand exchange where one strand of the donor molecule is displaced by the invading strand to form a D-loop. The invading strand is then extended using the homologous donor as a template [11, 12]. In the case of GC, only the area around the DSB is copied. GC may then proceed through a so-called synthesis-dependent strand annealing (SDSA) mechanism where the invading strand is displaced from the donor molecule and annealed to the sequences on the other end of the break (**Figure 1**). Alternatively, the second end of the break may be captured by the D-loop forming a double Holliday junction intermediate which can then be resolved by endonucleases or helicases (double-strand break repair pathway - DSBR) [11]. In contrast to SDSA, which generates exclusively non-crossovers, the resolution of a double Holliday junction may lead to either a crossover or a non-crossover. In BIR, the donor molecule is replicated from the invasion site all the way to the telomere by conservative replication (**Figure 1**) [15]. Both, GC and BIR may result in error-free repair if a sister chromatid is used as a donor. However, if the repair involves a homologous chromosome or an ectopic homology site, loss of heterozygosity or chromosomal rearrangements may occur [11, 12, 16].

The homology search and strand exchange reaction required for DSB repair by GC and BIR are catalysed by a recombinase called Rad51. Rad51 binds to resected ssDNA forming a Rad51-ssDNA filament (presynapsis) which then catalyses the search for homologous sequences and establishes a physical contact between the broken and donor molecules (synapsis) by invading the duplex donor DNA (**Figure 1**). As this step is absolutely required and is often rate-limiting in DSB repair, Rad51 assembly and disassembly plays an important role in modulating HR. In addition, disassembly of Rad51 filament is also required at the late stages of repair to restore the double-stranded structure of DNA [17]. Recently, a novel role for Rad51 in protection and restart of stalled replication forks has emerged [18]. Therefore, understanding the dynamics of Rad51 filaments is important for elucidating the molecular mechanisms of DNA repair and genome stability.

RAD51 RECOMBINASE

Rad51 was first genetically identified in 1974 in a screen for *Saccharomyces cerevisiae* mutants sensitive to ionising radiation [19]. Since then, the involvement of Rad51 in genome integrity and dynamics has been well-characterised, defining it as one of the key enzymes required for HR [20-25]. Rad51 is conserved among eukaryotes: Rad51 from vertebrates shares on average 74% protein sequence identity with fungi and plants while the human and mouse homologs are 99% identical [26]. Rad51 belongs to the ancient RecA/RAD51 protein family which apart from the bacterial and archaeal orthologues, RecA and RadA respectively, includes members that have diverged from the original function and adapted to more specialised roles. For example, Dmc1 is a meiosis-specific equivalent of Rad51, whereas yeast Rad55 and Rad57 are Rad51 paralogues involved in Rad51 regulation [27-30]. Although most eukaryotic orthologues are called Rad51 some, like the *Schizosaccharomyces pombe* orthologue Rhp51, may have a different name. For simplicity, these orthologues as a whole will be referred to as Rad51/RAD51 further in the review.

High protein sequence conservation suggests that the function of Rad51 is also conserved. Indeed, Rad51 deficient yeast and mouse trophoblast-like cells exhibit closely similar defects, including increased sensitivity to ionising radiation and chromosome loss, both of which can be explained by compromised DSB repair [19, 25, 31]. However, mammalian cells appear to be more sensitive to the loss of RAD51: a homozygous gene deletion results in embryonic lethality and failure to establish *Rad51* null cell lines [31]. This is also true for the chicken DT40 cells which accumulate spontaneous DNA DSBs in the absence of RAD51 and eventually die [32]. The stronger severity of the *RAD51* null phenotype in vertebrates compared to yeasts suggests a greater reliance of these cells on conserved RAD51 activities, possibly due to the emerging role of RAD51 in preventing DNA damage during conventional replication [18] or the higher frequency of stochastic DSBs per cell cycle stemming from the larger genome sizes [31].

Rad51 exists as a monomer in a solution but it can polymerise on both double-stranded DNA (dsDNA) and ssDNA in a cooperative manner and forms a right-handed nucleoprotein filament, in which the DNA is stretched and extended to facilitate homology search and base pairing [24, 33-37]. Unlike RecA, which has a very low affinity for dsDNA, Rad51 can bind both ssDNA and dsDNA *in vitro*, albeit with a preference for ssDNA [24, 38-40]. As required for its recombinase activity, Rad51 can bind two DNA molecules simultaneously, through its primary and secondary DNA binding sites. The primary binding site is responsible for the interactions with DNA during the formation of the filament and the double-stranded product in postsynapsis, while the secondary site is required for the capture of a donor molecule in presynapsis [41-43]. Rad51 can carry out the strand-exchange reaction *in vitro* (**Figure 2**). This activity is strongly stimulated by addition of the ssDNA binding protein RPA (Replication Protein A) after Rad51 is pre-




FIGURE 2: Rad51-catalysed strand exchange reaction. Circular ssDNA is pre-incubated with Rad51 to allow Rad51 binding to DNA without competition. RPA is then added to the reaction to remove secondary DNA structures. Nucleated Rad51 can replace RPA and form a functional continuous nucleofilament. When linear dsDNA is added Rad51 can catalyse the strand exchange between the double-stranded donor and the circular ssDNA.

nucleated with ssDNA [40, 44-48]. The positive effect of RPA was postulated to result from the ability of RPA to remove secondary DNA structures which might impede the formation of continuous, functional Rad51 filaments [47-49].

Rad51 also binds ATP and hydrolyses it in a DNA-dependent manner. The ATP binding is required for Rad51 activities [40, 44, 45, 50, 51]. Rad51 mutants deficient in nucleotide binding are catalytically dead and cannot form extended nucleofilaments, while mutants that bind ATP but are incapable of hydrolysing it can perform the strand exchange reaction *in vitro* and partially complement *RAD51* deletion *in vivo* when overexpressed [52, 53]. This demonstrates that only ATP binding is required for the essential Rad51 functions but ATP hydrolysis contributes to its full activity inside the cell. However, the role of Rad51 ATP hydrolysis is not completely understood. Possibly, it is important for the disassembly of the filament and recycling of Rad51 pool as *S. cerevisiae* Rad51, just like its bacterial orthologue RecA, shows decreased affinity for DNA when bound to ADP instead of ATP [54-56]. Consistent with this hypothesis, human RAD51 bound to DNA shows lower subunit turnover when ATP hydrolysis is prevented [37, 57]. Alternatively, the ATP hydrolysis might promote the DNA annealing step during the strand exchange reaction: while the ATP-bound nucleofilament conformation is stiffer and more suitable for the initial step of separating donor DNA strands, the ADP-bound human RAD51 might promote annealing of the invading and donor strands [58]. Interestingly, *in vitro* ATP hydrolysis by Rad51 is much slower than that by RecA. When bound to ssDNA, human and yeast orthologues hydrolyse 0.16 and 0.7 ATPs per minute per subunit respectively while RecA turns over 25-30 ATP molecules during that time [40, 44, 54, 59]. Therefore, the modulation of the ATPase activity of Rad51 *in vivo* by other proteins might have an important function in regulating Rad51 activity.

POSITIVE REGULATORS OF RAD51 NUCLEOPROTEIN FILAMENT FORMATION

As mentioned above, DSB repair by GC and BIR requires a formation of Rad51 filament on resected DNA. However, resected DNA is much more readily coated by the ssDNA

binding protein RPA, which is more abundant inside the cells and has a higher affinity for ssDNA than Rad51 [60]. As mentioned earlier, RPA stimulates Rad51-catalysed strand exchange *in vitro* when it is added after Rad51 has been assembled on ssDNA. However, if RPA is incubated with ssDNA before the Rad51 addition or both proteins are introduced into the reaction simultaneously, Rad51 filament formation and the subsequent strand exchange reaction are largely inhibited [46, 48, 61-63]. This indicates that Rad51 and RPA compete for binding to ssDNA and that Rad51 cannot efficiently replace RPA bound to ssDNA. However, the RPA/Rad51 protein exchange can be facilitated by so called mediator proteins described below.

Rad52 and BRCA2

Rad52 is a mediator which promotes Rad51 filament formation on RPA-coated ssDNA [46, 48, 62, 63]. Rad52 can physically interact with DNA, RPA and Rad51 and it has been postulated to promote the formation of Rad51 filaments by both recruiting Rad51 and helping it to replace RPA on ssDNA [46, 64-66]. Rad52 binds and stabilises RPA on ssDNA while promoting RPA replacement by Rad51 on the Rad52-surrounding DNA, thereby nucleating Rad51 filaments (Figure 3) [67]. The N-terminus of Rad51 has been demonstrated to interact with the DNA binding domain of RPA and possibly promote RPA dissociation [68]. This might allow the nucleated Rad51 to help free Rad51 monomers to compete for DNA binding, thereby extending the filament. This explanation is consistent with the recent *in vitro* reconstruction studies suggesting that RPA mainly inhibits the nucleation but not the elongation of Rad51 filaments [37, 67]. Rad52 also stabilises Rad51 filament via protein-protein interactions [69]. *S. cerevisiae* cells lacking Rad52 are deficient in DSB repair by HR and cannot form detectable DNA damage-induced Rad51 foci suggesting that Rad51 filament assembly requires Rad52 *in vivo* [70-72].

Rad52 orthologues are also present in vertebrates. However, RAD52 deficient mice, chicken and human cells show only a slight defect in HR and remain resistant to DSB inducing factors [73-75]. This is due to RAD52 playing a secondary role in recombination in mammals while another protein, BRCA2, mediates RAD51 filament formation [76,




FIGURE 3: A model of Rad51 nucleofilament formation in budding yeast. ssDNA generated as a result of resection or strand separation is rapidly covered by RPA (yellow spheres). Rad52 (purple spheres) binds RPA-coated ssDNA, recruits Rad51 (green spheres) and facilitates RPA-Rad51 exchange in the vicinity, thereby promoting Rad51-ssDNA filament formation. Rad55-Rad57 dimers (dark-green spheres) are also incorporated into the Rad51 nucleofilament and stabilise it by providing additional protein-protein interactions as well as antagonising Rad51 removal by Srs2 (teal ring). Only the main regulators of *S. cerevisiae* Rad51 are shown.

77]. Like *S. cerevisiae* Rad52, BRCA2 enables RAD51 to perform the strand exchange reaction with RPA-coated ssDNA substrate *in vitro* [77]. BRCA2 physically interacts with RAD51, RPA and DNA [78, 79]. BRCA2 has been proposed to stabilise RAD51 filaments *in vitro* by blocking ATP hydrolysis by RAD51 [77, 80]. In addition, one of the BRCA2-interacting proteins, BRCCIP β , also interacts with RAD51 and induces a conformational change which facilitates the normally slow release of ADP from RAD51, thereby pro-

moting nucleotide exchange and reversal of RAD51 to the active ATP-bound state [81, 82]. Another HR accessory factor, HOP2-MND1, can induce RAD51 conformational changes which enhance its ability to bind nucleotides and modulate the ssDNA-binding ability to promote the strand exchange reaction [83].

Interestingly, human RAD52 cannot substitute for BRCA2 in nucleating RAD51 on RPA-coated ssDNA *in vitro* but RAD52 depletion in BRCA2 deficient cells results in synthetic lethality accompanied by a further decrease in HR and a reduction in RAD51 foci induced by ionising radiation [77, 84]. This implies that although BRCA2 is the main RAD51 mediator in human cells, RAD52 activity in promoting the assembly of the RAD51 filament is significant enough to keep BRCA2-deficient cells viable.

Rad51 paralogues

Yeast Rad51 paralogues Rad55 and Rad57 are two other key mediators. They form a stable heterodimer which physically interacts with Rad51 and can alleviate the RPA-dependent inhibition of Rad51-catalysed strand exchange reaction *in vitro* [30]. The phenotype of the *rad55 rad57* double mutant is identical to those of individual deletions [85]. Mutants lacking Rad55 are sensitive to DNA-damaging agents, deficient in DSB repair by HR and show impaired localisation of Rad51 to an unreparable DSB suggesting that Rad51 filament assembly is defective in the absence of Rad55-Rad57 [70, 86]. The *rad51-I345T* mutation results in a Rad51 protein with increased affinity for DNA and the ability to overcome the inhibitory effect of RPA during the strand exchange reaction *in vitro*. *rad51-I345T* also partially suppresses the DNA damage sensitivity of *rad55 Δ* and *rad57 Δ* cells but it cannot compensate for the loss of the Rad52 mediator activity [85]. Rad55-Rad57 works downstream of the Rad52-dependent Rad51 recruitment to ssDNA and possibly stabilises Rad51 filament through protein-protein interactions, which might maximise the probability that a nucleation event will result in a successful assembly of a functional Rad51 nucleofilament. Rad55-Rad57 has been also proposed to enhance the stability of Rad51 filaments by antagonising Srs2 helicase (described below) [29] which dislodges Rad51 from ssDNA (Figure 3) [87, 88]. Consistently, Rad51-I345T which partially compensates for the loss of Rad55-Rad57 is harder for Srs2 to strip from ssDNA *in vitro* [89] and the DNA damage sensitivity of cells lacking the Rad55-Rad57 complex can be suppressed by a deletion of *SRS2* [87, 90, 91].

Five Rad51 paralogues have been described in vertebrates: RAD51B, RAD51C, RAD51D, XRCC2 and XRCC3 [27, 28]. They all function in the same pathway as BRCA2 and the depletion of any one of them results in a decreased efficiency of HR in human cells [92]. Phylogenetic studies suggest that XRCC2 and RAD51D are yeast Rad55 and Rad57 orthologues respectively, although some studies place Rad57 most closely related to XRCC3 [28, 93-95]. The Rad51 paralogs form two stable complexes: one is called BCDX2 and consists of RAD51B, RAD51C, RAD51D, and XRCC2 while the other one – CX3 – is made of RAD51C and XRCC3 [96]. The DNA damage induced formation of RAD51

foci is decreased in human cells depleted of RAD51D, RAD51C and XRCC2 but not XRCC3 suggesting that BCDX2 complex is required for efficient Rad51 filament formation, while CX3 is dispensable for this process [92, 97]. In agreement with this, the CX3 complex has been shown to act downstream of Rad51 filament assembly and participate in the resolution of Holliday junctions [98].

Shu complex

Another factor that participates in Rad51 filament assembly is the Shu complex. In *S. cerevisiae*, it consists of the Rad51 paralogues Csm2 and Psy3 which form a heterotetramer along with Shu1 and Shu2 [99, 100]. Unlike Rad52 and Rad55-Rad57, the Shu complex is not essential for DSB repair by HR *in vivo* [99]. However, deletions of the individual genes encoding the Shu complex components do lead to defects in Rad51 foci formation [100]. Csm2 physically interacts with Rad55 and bridges the Shu complex to Rad51 [101]. The Csm2-Psy3 dimer binds DNA and can enhance the Rad51-catalysed strand exchange reaction with RPA-coated substrates *in vitro*, in a Rad52- and Rad55-Rad57-dependent manner [101]. Although Shu1 and Shu2 are not required for this activity *in vitro*, *shu1Δ* results in elevated Srs2-dependent Rad51 filament disassembly *in vivo* [102]. Furthermore, Shu2 physically interacts with Srs2 suggesting that the Shu complex might promote Rad51 filament assembly by both direct interactions with Rad51 and inhibiting the Srs2-dependent disassembly of the filament [103].

A Shu2 orthologue – SWS1 – has been found in humans and shown to interact with another protein, SWSAP1. This interaction mutually stabilises the two proteins and the SWS1-SWSAP1 complex can bind ssDNA. SWSAP1 was also found to physically interact with RAD51, RAD51B, RAD51C, RAD51D and XRCC3 while SWS1 can bind to RAD51D and XRCC3 [104]. Depletion of either SWS1 or SWSAP1 results in decreased formation of DNA damage-induced RAD51 foci suggesting that the function of the Shu complex is conserved from yeast to humans [104, 105].

Rad54

Other factors, although less significant, have been implicated in promoting Rad51 nucleofilament formation. Rad54/RAD54 is an ATP-dependent translocase which can bind Rad51/RAD51 and DNA simultaneously and plays an important role in synaptic and postsynaptic events. While a catalytically-dead translocase is deficient in DNA strand invasion, it is fully functional in stabilising Rad51 filaments [106-108]. When Rad51 is assembled on dsDNA, Rad54 can strip it in an ATP-dependent manner [109, 110]. Therefore Rad54 promotes Rad51 filament formation on ssDNA by both stabilising Rad51 binding to ssDNA and inhibiting its association with dsDNA. However, cell cycle dependent phosphorylation of Rad54 can convert it into a negative regulator of Rad51, by enabling a Rad54-dependent removal of Rad51 at HR loci in G2 [111].

Swi5-Srf1 (Sae3-Mei5)

Sae3 and Mei5 are meiosis-specific mediators of Rad51 and Dmc1 filament formation in *S. cerevisiae* [112]. However, their conserved orthologues in other organisms, including *S. pombe*, mice and humans (Swi5/SWI5 and Sfr1/SFR1 respectively), participate in the assembly of mitotic Rad51/RAD51 filament [113-115]. Swi5/SWI5 and Sfr1/SFR1 form a stable complex with a 1:1 stoichiometry and can directly interact with Rad51/RAD51 [114-116]. *S. pombe* Swi5-Sfr1 stimulates the *in vitro* Rad51-dependent strand exchange reaction [117]. Furthermore, the formation of Rad51/RAD51 foci after exposure to ionising radiation is decreased in both *sfr1Δ S. pombe* and SRF1 depleted human cells, pointing towards a defect in the assembly of Rad51 [113, 114]. *S. pombe* Swi5 and Sfr1 proteins form an elongated structure *in vitro* which fits in the helical groove of the presynaptic filament, thereby suggesting a hypothetical mechanism for Swi5-Sfr1 action during the assembly of Rad51 [116].

INO80

It is important to note that Rad51 assembles on DNA in a context of chromatin. Not surprisingly, an evolutionally conserved nucleosome remodelling complex INO80 has been linked to the formation of the presynaptic filament [118-121]. Disruption of the INO80 remodelling complex in budding yeast and human cells results in a decreased efficiency of HR. Rad51 accumulation on resected DNA is also reduced pointing to a defect in the formation of the nucleofilament. This phenotype can be largely suppressed by a removal of the H2A.Z histone variant suggesting that H2A.Z has an inhibitory effect on the assembly of Rad51 and that INO80 complex is required to remove it from damaged chromatin [118, 119, 122, 123]. However, the exact mechanism of how H2A.Z inhibits Rad51 filament formation is unknown. Although DNA resection is also affected in cells lacking INO80 [118, 119], the H2A.Z removal suppresses the defect of Rad51 filament assembly but it does not compensate for the resection defect [119]. As HR efficiency is also largely restored in cells lacking both INO80 and H2A.Z, the main function of INO80 in HR is likely to be the facilitation of Rad51 nucleofilament formation [118, 119].

NEGATIVE REGULATION OF RAD51-DNA BINDING

Rad51 removal from undamaged dsDNA

As mentioned above, Rad51 can bind both ssDNA and dsDNA *in vitro* [24, 39]. While Rad51 binding to ssDNA is essential for the strand exchange reaction *in vitro*, pre-coating dsDNA with Rad51 inhibits the formation of the product [45, 109]. *In vivo*, Rad51 cellular pools are limited and Rad51 binding to dsDNA depletes the pool of free monomers leaving fewer of them available for the repair [70]. Thus, Rad51 binding to undamaged chromatin can impede Rad51-dependent recombination and may result in genome instability and chromosome loss [110, 124].

In *S. cerevisiae*, non-damage-associated DNA binding of Rad51 is cytologically undetectable as it is actively antagonised by three SWI2/SNF2 family DNA translocases – Rad54,

Rdh54 and Uls1 [124]. Both Rad54 and Rdh54 remove Rad51 from dsDNA in an ATP-dependent manner *in vitro* [109, 125]. This activity requires the N-terminal parts of Rad54 and Rdh54 which have been shown to interact with Rad51 but the exact mechanism of Rad51 displacement is unknown [126, 127]. Rad54 has been further demonstrated to enable the strand exchange reaction *in vitro*, even when both ssDNA and the donor dsDNA are covered with Rad51 [109].

Rdh54 might be the main player in the removal of Rad51 from undamaged chromatin as the lack of this protein but not the other two translocases results in accumulation of spontaneous non-damage-associated Rad51 foci [124]. However, deletion of all the three translocase-coding genes results in a more severe phenotype suggesting that Rad54 and Uls1 can partially substitute for Rdh54 [124, 128]. Consistent with the *in vitro* evidence, inactivation of the ATPase activity in any of the three proteins results in the same phenotype as in the corresponding deletion mutants, further highlighting the need for the Rdh54 and Rad54 translocase activities in the removal of Rad51 from dsDNA; Uls1 is likely to operate in a similar way [124].

Rad54 homologs are well-conserved among eukaryotes, with human cells containing two known proteins – RAD54 and RAD54B [129]. Like its yeast counterpart, human RAD54 can remove RAD51 from dsDNA *in vitro* [110]. Simultaneous depletion of both RAD54 and RAD54B results in accumulation of RAD51 on chromatin in human tumour cells [110]. The fact that both proteins need to be depleted to reveal the phenotype suggests that they are redundant and that RAD54B also may have the ability to remove RAD51 from dsDNA. This suggests that the function of the discussed SWI2/SNF2 translocases in the regulation of the RAD51 cellular pool might be conserved from yeasts to humans.

Suppression of HR at replication forks via inhibition of Rad51 filament formation

DNA synthesis in eukaryotes strongly depends on PCNA (Proliferating Cell Nuclear Antigen), a homotrimeric protein which forms a ring around the DNA and acts as a tool belt holding different components of the replication machinery: polymerases, ligases, nucleases, helicases, etc. [130]. Some of these enzymes, such as DNA polymerases, are associated with PCNA at the fork almost all the time while others can be recruited as needs for their activity at specific loci arise. For example, Pif1 family helicases are recruited to hard-to-replicated loci [131]. Replication fork barriers (tightly bound proteins, G-rich DNA regions, highly expressed genes, etc) cause replication fork pausing which often leads to accumulation of ssDNA, followed by recruitment of Rad52 and Rad51, thereby creating an opportunity for the DNA at the fork to be involved in unwanted HR. In budding yeast, these potentially mutagenic events are prevented by the Srs2 helicase which has been shown to disassemble the Rad51 nucleoprotein filament by dislodging Rad51 from ssDNA [87, 88]. Srs2 is recruited to replication forks through a direct binding of the SUMO-interacting motif (SIM) and the PCNA-interacting peptide

box in the C-terminus of Srs2 to a SUMOylated PCNA [132-134]. Once recruited to replication forks, Srs2 removes Rad51 and prevents potentially deleterious unscheduled HR events that might occur when replication fork progression is slowed down or paused by replication barriers. Disruption of this regulatory mechanism by mutations in *SRS2* leads to a hyper-recombination phenotype (increased mitotic recombination) [135, 136]. However, when replication forks are stalled due to a damaged template, Rad51 recruitment to the replication fork might be desirable in order to bypass the DNA lesion via the template switching mechanism [137]. In this case, local Srs2 levels might be decreased through targeting the fork-bound Srs2 for proteasomal degradation [138]. In addition, PCNA can be unloaded from stalled replication forks by the Elg1-containing RFC complex, thereby eliminating the PCNA-dependent binding of Srs2 to the fork [138].

Recent advances in understanding the role of RAD51 at replication forks in higher eukaryotes suggest an additional layer of regulatory mechanisms modulating RAD51 activities during replication. In human cells, a depletion of either BRCA2 or RAD51 results in under-replication and cell cycle arrest in the subsequent G1 phase [139]. The current understanding of replication considers stalled replication forks as part of normal cell physiology and fork reversal as a mechanism stabilising the forks [18]. RAD51 has been proposed to participate in both fork reversal [140] and protection of these forks from excessive DNA degradation by nucleases [141, 142]. A reversed replication fork is a four-way dsDNA junction, with one of the four branches terminating in a one-ended DSB which can serve as an entry point for the break resection machinery. BRCA2-dependent recruitment of RAD51 to a partially resected reversed fork has been proposed to inhibit further DNA degradation [142, 143]. On the other hand, RAD51 accumulation at replication forks is counteracted by a newly identified ssDNA-binding protein RADX which prevents fork collapse due to excessive activity of RAD51 [144]. Interestingly, deleting *RADX* restores fork protection in BRCA2-deficient cells [144] suggesting that a fine balance between the positive and negative regulation of RAD51 at the forks by BRCA2 and RADX respectively is required for the genome stability maintenance during replication. Rad51 was further demonstrated to physically interact with the primase Pol α , possibly by recruiting it to stalled replication forks and promoting their restart [142]. Rad51 also prevents the degradation of nascent DNA at ssDNA gaps which might form behind replication forks due to a damaged DNA template. Normally, these gaps are small and might be undetectable but accumulation of extensive ssDNA tracks behind replication forks has been observed in the absence of Rad51 binding to chromatin in *Xenopus* egg extracts. The accumulation of these ssDNA gaps may be suppressed by the inhibition of the Mre11 nuclease activity, further supporting the role of Rad51 in the protection of nascent DNA against the degradation by nucleases [141].

The role of yeast Rad51 in fork protection is less understood but some close similarities to the findings in higher eukaryotes have been found: *RAD51* deletion in

S. cerevisiae leads to accumulation of ssDNA gaps at forks and behind them. Rad51 and Rad52 localise to the forks during replication and are required for post-replicative DNA repair via HR [145]. Therefore, the replication-associated functions of Rad51 might be conserved in eukaryotes.

Disassembly of Rad51 nucleoprotein filament during DSB repair

On one hand, disassembly of Rad51 presynaptic filaments during DSB repair may play a role in limiting excessive recombination events. On the other hand, the same activity plays pro-recombination role as it is required at the late stages of repair to 'clean up' postsynaptic DNA in order to enable recruitment of PCNA and the rest of the replication machinery to re-synthesize resected DNA [17]. The Srs2 helicase has now been implicated in both functions and therefore Srs2-dependent disassembly of Rad51 filament previously considered inhibitory to HR, also has a pro-recombination role as discussed below.

PCNA is recruited to DNA by Replication Factor C (RFC) which recognises primer-template junctions and loads PCNA on dsDNA [146-148]. The RFC-PCNA complex can bind primer-template DNA junctions and in general has a significant affinity to ssDNA *in vitro*. The latter can be inhibited by the addition of RPA, which enhances the specificity of PCNA loading to the junctions [148-150]. RPA physically interacts with RFC and stimulates PCNA loading onto DNA *in vitro* [17, 151-153]. In contrast, Rad51 inhibits PCNA loading but this inhibition can be suppressed by either increasing the concentrations of RPA, which competes with Rad51 for ssDNA binding, or by addition of the Srs2 helicase which disassembles Rad51 filaments, thereby promoting RPA binding to DNA and the consequential PCNA recruitment via RPA-RFC interactions [17, 88, 152].

Srs2 can remove RPA, Rad52 and Rad51 from ssDNA *in vitro* [55, 87-89, 154]. Srs2 can physically interact with Rad51 and although *S. cerevisiae* Srs2 is capable of removing human RAD51 *in vitro*, the efficient clearance of ssDNA from Rad51 depends on species-specific interactions between Srs2 and Rad51, as well as ATP hydrolysis by Rad51 and Srs2 translocase activity [55, 89]. It has been suggested that Srs2 allosterically activates ATP hydrolysis in Rad51 monomers, thereby decreasing their affinity for DNA. Srs2 translocation is postulated to be important for the processivity and positioning of the helicase, which might be required to make appropriate contacts with the successive Rad51 monomers [55]. In addition, a tandem assembly of Srs2 monomers appears to be important for the efficient disassembly of Rad51 filaments [89]. On the other hand, the rate of Rad51 stripping is also influenced by the strength of the Rad51-ssDNA interaction: the amino acid substitutions which increase the stability of Rad51 on ssDNA due to either increased affinity for ssDNA or inability to hydrolyse ATP both slow down the rates of Rad51 removal by Srs2 [89].

Importantly, Srs2 has been proposed to promote SDSA by disrupting D-loops in a manner that requires its ATPase activity, Rad51-interacting domain, SIM and PCNA-

interacting peptide box [136, 155-157]. SRS2 deletion leads to an elevated frequency of crossovers during the repair of an induced DSB [155].

It is not known in detail how Srs2 is brought to DNA repair sites but its localisation is independent of the SIM which is required for Srs2 recruitment to stalled replication forks via interaction with SUMOylated PCNA [133]. The Srs2- ΔC (1-860) mutant protein lacking the Rad51-interacting region of Srs2 has a greatly impaired Rad51 clearance activity *in vitro*, mainly due to a decreased loading of mutant Srs2 on Rad51-ssDNA filaments [55, 89]. This suggests that the Rad51-interacting region of Srs2 promotes Srs2 association with the presynaptic filament [89]. However, Srs2 has been shown to localise to repair sites even in the absence of Rad51 [133]. Furthermore, the *srs2- ΔC* (1-860) allele can suppress the DNA re-synthesis defects arising from the lack of Rad51 removal in *srs2 Δ* cells [17]. These observations suggest that there might be a Rad51-independent way to recruit Srs2 to repair sites *in vivo*. Consistently, it has been recently demonstrated that Srs2 can directly bind heteroduplex DNA joints *in vitro* [89].

Inability to disassemble Rad51 filaments at the repair sites in *srs2 Δ* cells leads to accumulation of ssDNA gaps, persistence of the DNA damage signalling and failure to complete DNA repair [17, 158, 159]. These ssDNA gaps stem from a defect in re-synthesis of resected DNA which is likely required to terminate DNA resection. Although the re-synthesis is impaired in *srs2 Δ* , the reconstitution of dsDNA during DSB repair can be observed in *srs2 Δ* , albeit at a slower rate [17]. This could be explained by a sufficiently high stochastic exchange between Rad51 and RPA on ssDNA. Alternatively, there might be other yet unidentified proteins involved in Rad51 removal during the late stages of repair. Our recent experiments indicate that Rad54 might be such a protein as the loss of both Srs2 and Rad54 almost completely blocks re-synthesis of resected DNA during DSB repair (Andriuskevicius and Makovets, unpublished results). Our current hypothesis is that while Srs2 removes Rad51 from ssDNA, Rad54 might be disrupting the extension of the Rad51 filament at the neighbouring dsDNA region (**Figure 4**). In the absence of Srs2, the Rad54-dependent removal of Rad51 from dsDNA might extend into the ssDNA. Alternatively, stripping of Rad51 from dsDNA by Rad54 might promote stochastic dissociation of Rad51 from the ssDNA at the junction and vice versa: dislodging of Rad51 by Srs2 from ssDNA may destabilise Rad51 bound to the dsDNA at the junction (**Figure 4**). This hypothesis is consistent with the observations that stochastic Rad51 dissociation from DNA is higher for the monomers at the end of the filament than for the internal ones [37, 160]. The dissociation of Rad51 from the dsDNA at the junction might be required not only for the recruitment of PCNA but also to make the 3' end accessible for DNA polymerases in order to initiate DNA polymerisation, both during re-synthesis of resected DNA and when extending the invaded DNA strand in a D-loop [161]. This might explain why the post-invasion DNA synthesis during BIR in *srs2 Δ* is also affected [17]. To summarise, in yeast Rad54 and Srs2 might be complementing each other in disassembly of




FIGURE 4: A hypothetical model for the complementing roles of Rad54 and Srs2 in Rad51 removal at dsDNA-ssDNA junctions. In wild-type cells, Srs2 (teal ring) and Rad54 (orange ring) remove Rad51 from ssDNA and dsDNA respectively to allow RFC-PCNA (pink ellipsoid and light-brown ring) to access the dsDNA-ssDNA junction and recruit DNA polymerase (light-purple sphere). In the absence of Srs2, Rad54 removes Rad51 from dsDNA and either directly or indirectly promotes Rad51 dissociation from ssDNA at the junction.

Rad51 filaments while being partially redundant in Rad51 removal from the ssDNA-dsDNA junction due to the intrinsic features of the Rad51 filament. Indirect evidence suggest that RAD54 might have a similar function in vertebrates [111, 162].

A clear homolog of Srs2 has not been found in higher eukaryotes but other proteins have the ability to destabilise RAD51 nucleofilaments. G2-induced phosphorylation of Rad54 in *Xenopus* enables it to remove Rad51 from sites of HR [111]. Human helicases FBH1, BLM, RECQL5 and FANCD1 have been demonstrated to dislodge RAD51 from ssDNA in an ATP-dependent manner *in vitro* [163-166]. Another protein called PARI also has the ability to remove RAD51 from ssDNA *in vitro*. Although PARI does not have an ATPase activity, the PARI-mediated disassembly of the nucleofilament is dependent on ATP hydrolysis by RAD51 suggesting that PARI, like Srs2, can stimulate RAD51 ATPase activity [167]. This also supports the idea that ATP hydrolysis destabilises human RAD51 filament. Similarly to Srs2, PARI also functions at replication forks where it is recruited via PCNA and PCNA SUMOylation facilitates this recruitment [167]. Mammalian cells lacking FBH1, BLM, RECQL5 or PARI exhibit hyper-recombination phenotypes suggesting the relevance of these proteins to the negative control of recombination *in vivo* [165, 167-170].

Human cells might also be able to remove RAD51 from DNA at DNA damage sites via ubiquitination [162]. The RFW3 E3 ubiquitin ligase can physically interact with

RAD51 and polyubiquitinate it both *in vitro* and *in vivo*. The RFW3-dependent ubiquitination of RAD51 results in its degradation by proteasome. Cells depleted of RFW3 or expressing an ubiquitination-deficient RAD51 variant show decreased turnover of RAD51 at repair sites, suggesting that the ubiquitination might destabilise RAD51 in the filament by targeting it for degradation. Although the exact stage of HR involving RFW3-dependent regulation is unknown, it is proposed to function after the formation of RAD51 nucleofilament, and perhaps, even downstream of postsynapsis. If RFW3 were destabilising RAD51 at the presynaptic stage and working as an anti-recombinase, the HR frequency would have been increased when RAD51 ubiquitination was prevented. However, the opposite has been observed in cells expressing ubiquitination-defective RAD51 mutant protein, suggesting that RFW3-assisted RAD51 removal is important for the progression of HR at the later stages, possibly during the postsynaptic strand extension [162]. Interestingly, decreased RAD54 chromatin loading was observed in RFW3 deficient human cells after DNA damage, suggesting that RAD51 ubiquitination and RAD54 loading might be functionally linked [162].

It is important to emphasise that Rad51 filament is formed on resected DNA breaks independently of whether the repair will proceed by HR or by some other mechanism, for example, SSA or DNTA. Whichever route the repair takes, eventually the resected DNA is to be re-synthesised and converted back to dsDNA, in order to complete the

repair and switch off the DNA damage signalling. Although Rad51 is not required for DSB repair via either SSA or DNTA, the Srs2-dependent activity on Rad51 is necessary for these mechanisms to be efficient [17]. The results on DNTA are particularly interesting: they suggest that DSB healing by telomerase normally happens on resected DNA ends, although DNTA is dramatically increased in mutants with deficient resection [171, 172]. The evidence for DNTA and BIR raises a possibility that any DNA repair mechanisms that involve generation of ssDNA might be affected by unwanted Rad51 filament formation if the ssDNA is persistent long enough for the filament to assemble. Therefore, Rad51 removal by Srs2 or other means might be required not only during DSB repair but have a broader function in DNA repair.

REGULATION OF RAD51 FILAMENT DYNAMICS THROUGH POST-TRANSLATIONAL MODIFICATIONS

Rad51 functions are regulated through post-translational modifications (PTMs) of not only Rad51 but also of the positive and negative regulators of Rad51 nucleoprotein filament assembly. The purpose of most of these modifications falls into one of the three categories: i) cell cycle dependent restrictions on the activity of a factor; ii) upregulation of DNA repair in cells with DNA damage; and iii) regulation of protein turnover at repair sites. Homology-based repair is limited to S-G2, mostly due to the cell cycle dependent phosphorylation required to activate the resection endonucleases upon entry in the S phase. This might explain why Rad52 localisation to DNA depends on the CDK1 activity [173, 174] although CDK1-dependent phosphorylation of Rad52 has been detected in high-throughput experiments [175] and therefore Rad52 might be also directly regulated by CDK1. SUMOylation is involved in regulation of Rad52 too: it leads to stabilisation of the protein *in vivo* [176] while decreasing its affinity to ssDNA [177]. In yeast, both Rad51 and Rad55 are phosphorylated in response to DNA damage and these PTMs are required for cell survival upon DNA damage induction [178-180]. Activation of the DNA damage response in yeast as well as mammals leads to RPA SUMOylation [181-183] which increases its interaction with RAD51, possibly through the recently identified SIM within RAD51 [184]. Both RAD51 and BRCA2 are deubiquitinated in response to DNA damage [185, 186]: deubiquitination of BRCA2 leads to its stabilisation [186] while deubiquitinated RAD51 increases its binding to BRCA2 [185]. In turn, RAD51 ubiquitination might be important for RAD51 removal at the later stages of repair possibly by RAD54 [162], as discussed above. The removal of Rad51 from repair foci in G2 in *Xenopus* is promoted by cell cycle specific phosphorylation of Rad54 [111].

The role of multiple PTMs in regulation of Srs2 is well-documented. CDK1-dependent phosphorylation at the C-terminus promotes the interaction of Srs2 with Mre11 [187], counteracts its SUMOylation and directs Srs2 to unwinding D-loops, thereby promoting DSB repair by SDSA [188]. Although this phosphorylation controls the localisa-

tion of the Srs2 to D-loops, neither the elimination of the phosphorylation nor *SRS2* deletion has an effect on Rad51 presence at the site of strand invasion, suggesting that Srs2 acts downstream of the invasion step [188]. The C-terminus of Srs2 also plays an important role in regulation of its function at replication forks. As mentioned above, the C-terminus of Srs2 contains a SIM which is required for Srs2 interaction with PCNA at replication forks. However, the C-terminus also contains SUMOylation sites and Srs2 SUMOylation inhibits its interaction with PCNA [189]. Therefore, PTMs of Srs2 regulate both its anti-recombinase role at replication forks and its pro-recombination function in SDSA. The C-terminus is not required for the role of Srs2 in Rad51 removal during re-synthesis of resected DNA [17]. However, the shortest functional C-terminal truncation Srs2 (1-860) retains two out of the seven identified CDK1 sites which might be important for this function. In summary, regulation of HR in general and Rad51 filament assembly in particular through PTM provides an additional layer of mechanisms to boost Rad51 activities in response to DNA damage, fine-tune them depending on the type of DNA damage and cell cycle stage as well as enables cells to regulate Rad51 and its interaction with DNA and other proteins as the repair progresses.

CONCLUSION

The importance of recombinases in genome stability and diversity has been appreciated since the early days of molecular biology. Classical biochemistry and crystallography followed by more recently emerged single-molecule analysis and cryo-electron microscopy have provided molecular insights into their structure and function at the molecular level. We have also learned that numerous accessory proteins are required for a recombinase to operate *in vivo*; a whole set of enzymes built around it comprises HR machinery. The complexity of regulatory mechanisms governing this machinery has been becoming more and more obvious with the increasing number of discoveries of PTMs which modulate the components of HR machinery and their regulators. Rad51 filament formation is one of the most critical steps in HR and its regulation, both positive (filament assembly) and negative (disassembly) are required for efficient DNA repair. The newly-emerging role of recombinases at stalled replication forks adds another angle to understanding biological significance of their regulation. The sophistication of this regulatory network, based on the multitude of inputs and connections to other cellular pathways, increases from prokaryotes to yeast to higher eukaryotes, thereby underlying the importance of the fine-tuning of recombinase activities for cell survival and adaptive evolution of species.

ACKNOWLEDGMENTS

We are grateful to Benura Azeroglu, Elizabeth Bayne and Sara Buonomo for critical reading of the manuscript and helpful suggestions. T.A. thanks BBSRC and the School of Biological Sciences, University of Edinburgh for an EastBio doctoral training partnership stipend (grant number

BB/M010996/1). O.K. is a recipient of an Edinburgh Global Research Scholarship from the School of Biological Sciences. The Makovets lab is funded by the Leverhulme Trust Project Grant RPG-2016-379 to S.M.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

COPYRIGHT

© 2018 Andriuskevicius *et al.* This is an open-access article released under the terms of the Creative Commons Attribution

license (CC BY) license, which allows the unrestricted use, distribution, and reproduction in any medium, provided the original author and source are acknowledged.

Please cite this article as: Tadas Andriuskevicius, Oleksii Kottenko and Svetlana Makovets (2018). Putting together and taking apart: assembly and disassembly of the Rad51 nucleoprotein filament in DNA repair and genome stability. *Cell Stress* 2(5): 96-112. doi: 10.15698/cst2018.05.134

REFERENCES

- McKinnon PJ, Caldecott KW (2007). DNA strand break repair and human genetic disease. *Annual review of genomics and human genetics* 8:37-55. doi: 10.1146/annurev.genom.7.080505.115648
- Rodgers K, McVey M (2016). Error-prone repair of DNA double-strand breaks. *J Cell Physiol* 231(1): 15-24. doi: 10.1002/jcp.25053
- Kramer KM, Brock JA, Bloom K, Moore JK, Haber JE (1994). Two different types of double-strand breaks in *Saccharomyces cerevisiae* are repaired by similar RAD52-independent, nonhomologous recombination events. *Mol Cell Biol* 14(2): 1293-1301. doi: 10.1128/mcb.14.2.1293
- Gorbunova V, Levy AA (1997). Non-homologous DNA end joining in plant cells is associated with deletions and filler DNA insertions. *Nucleic Acids Res* 25(22): 4650-4657. doi: 10.1093/nar/25.22.4650
- Smogorzewska A, Karlseder J, Holtgreve-Grez H, Jauch A, de Lange T (2002). DNA ligase IV-dependent NHEJ of deprotected mammalian telomeres in G1 and G2. *Curr Biol* 12(19): 1635-1644. doi: 10.1016/s0960-9822(02)01179-x
- Ghezraoui H, Piganeau M, Renouf B, Renaud JB, Sallmyr A, Ruis B, Oh S, Tomkinson AE, Hendrickson EA, Giovannangeli C, Jasin M, Brunet E (2014). Chromosomal translocations in human cells are generated by canonical nonhomologous end-joining. *Mol Cell* 55(6): 829-842. doi: 10.1016/j.molcel.2014.08.002
- Zamborszky J, Szikriszt B, Gervai JZ, Pipek O, Poti A, Krzystanek M, Ribli D, Szalai-Gindl JM, Csabai I, Szallasi Z, Swanton C, Richardson AL, Szuts D (2017). Loss of BRCA1 or BRCA2 markedly increases the rate of base substitution mutagenesis and has distinct effects on genomic deletions. *Oncogene* 36(6): 746-755. doi: 10.1038/ncr.2016.243
- Jackson SP (2002). Sensing and repairing DNA double-strand breaks. *Carcinogenesis* 23(5): 687-696. doi: 10.1093/carcin/23.5.687
- Bee L, Fabris S, Cherubini R, Mognato M, Celotti L (2013). The efficiency of homologous recombination and non-homologous end joining systems in repairing double-strand breaks during cell cycle progression. *PLoS One* 8(7): e69061. doi: 10.1371/journal.pone.0069061
- Ribeyre C, Shore D (2013). Regulation of telomere addition at DNA double-strand breaks. *Chromosoma* 122(3): 159-173. doi: 10.1007/s00412-013-0404-2
- Chen JM, Cooper DN, Chuzhanova N, Ferec C, Patrinos GP (2007). Gene conversion: mechanisms, evolution and human disease. *Nat Rev Genet* 8(10): 762-775. doi: 10.1038/nrg2193
- Malkova A, Ira G (2013). Break-induced replication: functions and molecular mechanism. *Curr Opin Genet Dev* 23(3): 271-279. doi: 10.1016/j.gde.2013.05.007
- Bhargava R, Onyango DO, Stark JM (2016). Regulation of Single-Strand Annealing and its Role in Genome Maintenance. *Trends Genet* 32(9): 566-575. doi: 10.1016/j.tig.2016.06.007
- Symington LS (2016). Mechanism and regulation of DNA end resection in eukaryotes. *Crit Rev Biochem Mol Biol* 51(3): 195-212. doi: 10.3109/10409238.2016.1172552
- Saini N, Ramakrishnan S, Elango R, Ayyar S, Zhang Y, Deem A, Ira G, Haber JE, Lobachev KS, Malkova A (2013). Migrating bubble during break-induced replication drives conservative DNA synthesis. *Nature* 502(7471): 389-392. doi: 10.1038/nature12584
- Stark JM, Jasin M (2003). Extensive loss of heterozygosity is suppressed during homologous repair of chromosomal breaks. *Mol Cell Biol* 23(2): 733-743. doi: 10.1128/mcb.23.2.733-743.2003
- Vasianovich Y, Altmannova V, Kottenko O, Newton MD, Krejci L, Makovets S (2017). Unloading of homologous recombination factors is required for restoring double-stranded DNA at damage repair loci. *EMBO J* 36(2): 213-231. doi: 10.15252/embj.201694628
- Kolinjivadi AM, Sannino V, de Antoni A, Techer H, Baldi G, Costanzo V (2017). Moonlighting at replication forks - a new life for homologous recombination proteins BRCA1, BRCA2 and RAD51. *FEBS letters* 591(8): 1083-1100. doi: 10.1002/1873-3468.12556
- Game JC, Mortimer RK (1974). A genetic study of X-ray sensitive mutants in yeast. *Mutat Res* 24(3): 281-292. doi: 10.1016/0165-1218(74)90185-2
- Nakai S, Matsumoto S (1967). Two types of radiation-sensitive mutant in yeast. *Mutat Res* 4(2): 129-136. doi: 10.1016/0027-5107(67)90064-4
- Hama-Inaba H, Saeki T (1975). Evidence that a recombinationless strain, rad 51, of *Saccharomyces cerevisiae* lacks the budding cell resistance to gamma-rays. *J Radiat Res* 16(1): 37-44. doi: 10.1269/jrr.16.37
- Resnick MA (1976). The repair of double-strand breaks in DNA; a model involving recombination. *J Theor Biol* 59(1): 97-106. doi: 10.1016/s0022-5193(76)80025-2
- Jachymczyk WJ, von Borstel RC, Mowat MR, Hastings PJ (1981). Repair of interstrand cross-links in DNA of *Saccharomyces cerevisiae* requires two systems for DNA repair: the RAD3 system and the RAD51 system. *Mol Gen Genet* 182(2): 196-205. doi: 10.1007/bf00269658
- Shinohara A, Ogawa H, Ogawa T (1992). Rad51 protein involved in repair and recombination in *S. cerevisiae* is a RecA-like protein. *Cell* 69(3): 457-470. doi: 10.1016/0092-8674(92)90447-k
- Malkova A, Ivanov EL, Haber JE (1996). Double-strand break repair in the absence of RAD51 in yeast: a possible role for break-induced DNA replication. *Proc Natl Acad Sci U S A* 93(14): 7131-7136. doi: 10.1073/pnas.93.14.7131

26. Stassen NY, Logsdon JM, Jr., Vora GJ, Offenbergh HH, Palmer JD, Zolan ME (1997). Isolation and characterization of rad51 orthologs from *Coprinus cinereus* and *Lycopersicon esculentum*, and phylogenetic analysis of eukaryotic recA homologs. *Curr Genet* 31(2): 144-157. doi: 10.1007/s002940050189
27. Chintapalli SV, Bhardwaj G, Babu J, Hadjiyianni L, Hong Y, Todd GK, Boosalis CA, Zhang Z, Zhou X, Ma H, Anishkin A, van Rossum DB, Patterson RL (2013). Reevaluation of the evolutionary events within recA/RAD51 phylogeny. *BMC Genomics* 14(240). doi: 10.1186/1471-2164-14-240
28. Lin Z, Kong H, Nei M, Ma H (2006). Origins and evolution of the recA/RAD51 gene family: evidence for ancient gene duplication and endosymbiotic gene transfer. *Proc Natl Acad Sci U S A* 103(27): 10328-10333. doi: 10.1073/pnas.0604232103
29. Liu J, Renault L, Veaute X, Fabre F, Stahlberg H, Heyer WD (2011). Rad51 paralogues Rad55-Rad57 balance the antirecombinase Srs2 in Rad51 filament formation. *Nature* 479(7372): 245-248. doi: 10.1038/nature10522
30. Sung P (1997). Yeast Rad55 and Rad57 proteins form a heterodimer that functions with replication protein A to promote DNA strand exchange by Rad51 recombinase. *Genes Dev* 11(9): 1111-1121. doi: 10.1101/gad.11.9.1111
31. Lim DS, Hasty P (1996). A mutation in mouse rad51 results in an early embryonic lethal that is suppressed by a mutation in p53. *Mol Cell Biol* 16(12): 7133-7143. doi: 10.1128/mcb.16.12.7133
32. Sonoda E, Sasaki MS, Buerstedde JM, Bezzubova O, Shinohara A, Ogawa H, Takata M, Yamaguchi-Iwai Y, Takeda S (1998). Rad51-deficient vertebrate cells accumulate chromosomal breaks prior to cell death. *EMBO J* 17(2): 598-608. doi: 10.1093/emboj/17.2.598
33. Yu X, Jacobs SA, West SC, Ogawa T, Egelman EH (2001). Domain structure and dynamics in the helical filaments formed by RecA and Rad51 on DNA. *Proc Natl Acad Sci USA* 98(15): 8419-8424. doi: 10.1073/pnas.111005398
34. Danilowicz C, Peacock-Villada A, Vlassakis J, Facon A, Feinstein E, Kleckner N, Prentiss M (2014). The differential extension in dsDNA bound to Rad51 filaments may play important roles in homology recognition and strand exchange. *Nucleic Acids Res* 42(1): 526-533. doi: 10.1093/nar/gkt867
35. Short JM, Liu Y, Chen S, Soni N, Madhusudhan MS, Shivji MK, Venkataraman AR (2016). High-resolution structure of the presynaptic RAD51 filament on single-stranded DNA by electron cryo-microscopy. *Nucleic Acids Res* 44(19): 9017-9030. doi: 10.1093/nar/gkw783
36. Conway AB, Lynch TW, Zhang Y, Fortin GS, Fung CW, Symington LS, Rice PA (2004). Crystal structure of a Rad51 filament. *Nat Struct Mol Biol* 11(8): 791-796. doi: 10.1038/nsmb795
37. Ma CJ, Gibb B, Kwon Y, Sung P, Greene EC (2017). Protein dynamics of human RPA and RAD51 on ssDNA during assembly and disassembly of the RAD51 filament. *Nucleic Acids Res* 45(2): 749-761. doi: 10.1093/nar/gkw1125
38. McEntee K, Weinstock GM, Lehman IR (1981). Binding of the recA protein of *Escherichia coli* to single- and double-stranded DNA. *J Biol Chem* 256(16): 8835-8844. PMID: 7021553
39. Zaitseva EM, Zaitsev EN, Kowalczykowski SC (1999). The DNA binding properties of *Saccharomyces cerevisiae* Rad51 protein. *J Biol Chem* 274(5): 2907-2915. PMID: 9915828
40. Baumann P, Benson FE, West SC (1996). Human Rad51 protein promotes ATP-dependent homologous pairing and strand transfer reactions in vitro. *Cell* 87(4): 757-766. doi: 10.1016/s0092-8674(00)81394-x
41. Prasad TK, Yeykal CC, Greene EC (2006). Visualizing the assembly of human Rad51 filaments on double-stranded DNA. *J Mol Biol* 363(3): 713-728. doi: 10.1016/j.jmb.2006.08.046
42. Zhang XP, Galkin VE, Yu X, Egelman EH, Heyer WD (2009). Loop 2 in *Saccharomyces cerevisiae* Rad51 protein regulates filament formation and ATPase activity. *Nucleic Acids Res* 37(1): 158-171. doi: 10.1093/nar/gkn914
43. Matsuo Y, Sakane I, Takizawa Y, Takahashi M, Kurumizaka H (2006). Roles of the human Rad51 L1 and L2 loops in DNA binding. *FEBS J* 273(14): 3148-3159. doi: 10.1111/j.1742-4658.2006.05323.x
44. Sung P (1994). Catalysis of ATP-dependent homologous DNA pairing and strand exchange by yeast RAD51 protein. *Science* 265(5176): 1241-1243. doi: 10.1126/science.8066464
45. Sung P, Robberson DL (1995). DNA strand exchange mediated by a RAD51-ssDNA nucleoprotein filament with polarity opposite to that of RecA. *Cell* 82(3): 453-461. doi: 10.1016/0092-8674(95)90434-4
46. Sung P (1997). Function of yeast Rad52 protein as a mediator between replication protein A and the Rad51 recombinase. *J Biol Chem* 272(45): 28194-28197. doi: 10.1074/jbc.272.45.28194
47. Sugiyama T, Zaitseva EM, Kowalczykowski SC (1997). A single-stranded DNA-binding protein is needed for efficient presynaptic complex formation by the *Saccharomyces cerevisiae* Rad51 protein. *J Biol Chem* 272(12): 7940-7945. doi: 10.1074/jbc.272.12.7940
48. Song B, Sung P (2000). Functional interactions among yeast Rad51 recombinase, Rad52 mediator, and replication protein A in DNA strand exchange. *J Biol Chem* 275(21): 15895-15904. doi: 10.1074/jbc.M910244199
49. Eggler AL, Inman RB, Cox MM (2002). The Rad51-dependent pairing of long DNA substrates is stabilized by replication protein A. *J Biol Chem* 277(42): 39280-39288. doi: 10.1074/jbc.M204328200
50. Ogawa T, Yu X, Shinohara A, Egelman EH (1993). Similarity of the yeast RAD51 filament to the bacterial RecA filament. *Science* 259(5103): 1896-1899. doi: 10.1126/science.8456314
51. Benson FE, Stasiak A, West SC (1994). Purification and characterization of the human Rad51 protein, an analogue of *E. coli* RecA. *EMBO J* 13(23): 5764-5771. PMID: 7988572
52. Morrison C, Shinohara A, Sonoda E, Yamaguchi-Iwai Y, Takata M, Weichselbaum RR, Takeda S (1999). The essential functions of human Rad51 are independent of ATP hydrolysis. *Mol Cell Biol* 19(10): 6891-6897. doi: 10.1128/mcb.19.10.6891
53. Sung P, Stratton SA (1996). Yeast Rad51 recombinase mediates polar DNA strand exchange in the absence of ATP hydrolysis. *J Biol Chem* 271(45): 27983-27986. doi: 10.1074/jbc.271.45.27983
54. Menetski JP, Varghese A, Kowalczykowski SC (1988). Properties of the high-affinity single-stranded DNA binding state of the *Escherichia coli* recA protein. *Biochemistry* 27(4): 1205-1212. doi: 10.1021/bi00404a021
55. Antony E, Tomko EJ, Xiao Q, Krejci L, Lohman TM, Ellenberger T (2009). Srs2 disassembles Rad51 filaments by a protein-protein interaction triggering ATP turnover and dissociation of Rad51 from DNA. *Mol Cell* 35(1): 105-115. doi: 10.1016/j.molcel.2009.05.026
56. Shim KS, Schmutte C, Yoder K, Fishel R (2006). Defining the salt effect on human RAD51 activities. *DNA Repair (Amst)* 5(6): 718-730. doi: 10.1016/j.dnarep.2006.03.006
57. Chi P, Van Komen S, Sehorn MG, Sigurdsson S, Sung P (2006). Roles of ATP binding and ATP hydrolysis in human Rad51 recombinase function. *DNA Repair* 5(3): 381-391. doi: 10.1016/j.dnarep.2005.11.005

58. Kim HK, Morimatsu K, Norden B, Ardhammar M, Takahashi M (2002). ADP stabilizes the human Rad51-single stranded DNA complex and promotes its DNA annealing activity. **Genes Cells** 7(11): 1125-1134. doi: 10.1046/j.1365-2443.2002.00588.x
59. Brenner SL, Mitchell RS, Morrical SW, Neuendorf SK, Schutte BC, Cox MM (1987). recA protein-promoted ATP hydrolysis occurs throughout recA nucleoprotein filaments. **J Biol Chem** 262(9): 4011-4016. PMID: 2951381
60. Wang X, Haber JE (2004). Role of *Saccharomyces* single-stranded DNA-binding protein RPA in the strand invasion step of double-strand break repair. **PLoS Biol** 2(1): E21. doi: 10.1371/journal.pbio.0020021
61. Sugiyama T, Kowalczykowski SC (2002). Rad52 protein associates with replication protein A (RPA)-single-stranded DNA to accelerate Rad51-mediated displacement of RPA and presynaptic complex formation. **J Biol Chem** 277(35): 31663-31672. doi: 10.1074/jbc.M203494200
62. Shinohara A, Ogawa T (1998). Stimulation by Rad52 of yeast Rad51-mediated recombination. **Nature** 391(6665): 404-407. doi: 10.1038/34943
63. New JH, Sugiyama T, Zaitseva E, Kowalczykowski SC (1998). Rad52 protein stimulates DNA strand exchange by Rad51 and replication protein A. **Nature** 391(6665): 407-410. doi: 10.1038/34950
64. Milne GT, Weaver DT (1993). Dominant negative alleles of RAD52 reveal a DNA repair/recombination complex including Rad51 and Rad52. **Genes Dev** 7(9): 1755-1765. doi: 10.1101/gad.7.9.1755
65. Shinohara A, Shinohara M, Ohta T, Matsuda S, Ogawa T (1998). Rad52 forms ring structures and co-operates with RPA in single-strand DNA annealing. **Genes Cells** 3(3): 145-156. doi: 10.1046/j.1365-2443.1998.00176.x
66. Hays SL, Firmenich AA, Massey P, Banerjee R, Berg P (1998). Studies of the interaction between Rad52 protein and the yeast single-stranded DNA binding protein RPA. **Mol Cell Biol** 18(7): 4400-4406. doi: 10.1128/mcb.18.7.4400
67. Gibb B, Ye LF, Kwon Y, Niu H, Sung P, Greene EC (2014). Protein dynamics during presynaptic-complex assembly on individual single-stranded DNA molecules. **Nat Struct Mol Biol** 21(10): 893-900. doi: 10.1038/nsmb.2886
68. Stauffer ME, Chazin WJ (2004). Physical interaction between replication protein A and Rad51 promotes exchange on single-stranded DNA. **J Biol Chem** 279(24): 25638-25645. doi: 10.1074/jbc.M400029200
69. New JH, Kowalczykowski SC (2002). Rad52 protein has a second stimulatory role in DNA strand exchange that complements replication protein-A function. **J Biol Chem** 277(29): 26171-26176. doi: 10.1074/jbc.M203670200
70. Sugawara N, Wang X, Haber JE (2003). In vivo roles of Rad52, Rad54, and Rad55 proteins in Rad51-mediated recombination. **Mol Cell** 12(1): 209-219. doi: 10.1016/s1097-2765(03)00269-7
71. Miyazaki T, Bressan DA, Shinohara M, Haber JE, Shinohara A (2004). In vivo assembly and disassembly of Rad51 and Rad52 complexes during double-strand break repair. **EMBO J** 23(4): 939-949. doi: 10.1038/sj.emboj.7600091
72. Lisby M, Barlow JH, Burgess RC, Rothstein R (2004). Choreography of the DNA damage response: spatiotemporal relationships among checkpoint and repair proteins. **Cell** 118(6): 699-713. doi: 10.1016/j.cell.2004.08.015
73. Rijkers T, Van Den Ouweland J, Morolli B, Rolink AG, Baarends WM, Van Sloun PP, Lohman PH, Pastink A (1998). Targeted inactivation of mouse RAD52 reduces homologous recombination but not resistance to ionizing radiation. **Mol Cell Biol** 18(11): 6423-6429. doi: 10.1128/mcb.18.11.6423
74. Yamaguchi-Iwai Y, Sonoda E, Buerstedde JM, Bezzubova O, Morrison C, Takata M, Shinohara A, Takeda S (1998). Homologous recombination, but not DNA repair, is reduced in vertebrate cells deficient in RAD52. **Mol Cell Biol** 18(11): 6430-6435. doi: 10.1128/mcb.18.11.6430
75. Kan Y, Batada NN, Hendrickson EA (2017). Human somatic cells deficient for RAD52 are impaired for viral integration and compromised for most aspects of homology-directed repair. **DNA Repair** 55:64-75. doi: 10.1016/j.dnarep.2017.04.006
76. Davies AA, Masson JY, Mcllwraith MJ, Stasiak AZ, Stasiak A, Venkitaraman AR, West SC (2001). Role of BRCA2 in control of the RAD51 recombination and DNA repair protein. **Mol Cell** 7(2): 273-282. doi: 10.1016/s1097-2765(01)00175-7
77. Jensen RB, Carreira A, Kowalczykowski SC (2010). Purified human BRCA2 stimulates RAD51-mediated recombination. **Nature** 467(7316): 678-683. doi: 10.1038/nature09399
78. Scully R, Chen J, Plug A, Xiao Y, Weaver D, Feunteun J, Ashley T, Livingston DM (1997). Association of BRCA1 with Rad51 in mitotic and meiotic cells. **Cell** 88(2): 265-275. doi: 10.1016/s0092-8674(00)81847-4
79. Wong JM, Ionescu D, Ingles CJ (2003). Interaction between BRCA2 and replication protein A is compromised by a cancer-predisposing mutation in BRCA2. **Oncogene** 22(1): 28-33. doi: 10.1038/sj.onc.1206071
80. Carreira A, Hilarion J, Amitani I, Baskin RJ, Shivji MK, Venkitaraman AR, Kowalczykowski SC (2009). The BRC repeats of BRCA2 modulate the DNA-binding selectivity of RAD51. **Cell** 136(6): 1032-1043. doi: 10.1016/j.cell.2009.02.019
81. Kelso AA, Goodson SD, Watts LE, Ledford LL, Waldvogel SM, Diehl JN, Shah SB, Say AF, White JD, Sehorn MG (2017). The beta-isoform of BCCIP promotes ADP release from the RAD51 presynaptic filament and enhances homologous DNA pairing. **Nucleic Acids Res** 45(2): 711-725. doi: 10.1093/nar/gkw877
82. Bugreev DV, Mazin AV (2004). Ca²⁺ activates human homologous recombination protein Rad51 by modulating its ATPase activity. **Proc Natl Acad Sci U S A** 101(27): 9988-9993. doi: 10.1073/pnas.0402105101
83. Bugreev DV, Huang F, Mazina OM, Pezza RJ, Voloshin ON, Camerini-Otero RD, Mazin AV (2014). HOP2-MND1 modulates RAD51 binding to nucleotides and DNA. **Nat Commun** 5:4198. doi: 10.1038/ncomms5198
84. Feng Z, Scott SP, Bussen W, Sharma GG, Guo G, Pandita TK, Powell SN (2011). Rad52 inactivation is synthetically lethal with BRCA2 deficiency. **Proc Natl Acad Sci U S A** 108(2): 686-691. doi: 10.1073/pnas.1010959107
85. Fortin GS, Symington LS (2002). Mutations in yeast Rad51 that partially bypass the requirement for Rad55 and Rad57 in DNA repair by increasing the stability of Rad51-DNA complexes. **EMBO J** 21(12): 3160-3170. doi: 10.1093/emboj/cdf293
86. Hays SL, Firmenich AA, Berg P (1995). Complex formation in yeast double-strand break repair: participation of Rad51, Rad52, Rad55, and Rad57 proteins. **Proc Natl Acad Sci USA** 92(15): 6925-6929. doi: 10.1073/pnas.92.15.6925
87. Veaute X, Jeusset J, Soustelle C, Kowalczykowski SC, Le Cam E, Fabre F (2003). The Srs2 helicase prevents recombination by disrupting Rad51 nucleoprotein filaments. **Nature** 423(6937): 309-312. doi: 10.1038/nature01585

88. Krejci L, Van Komen S, Li Y, Villemain J, Reddy MS, Klein H, Ellenberger T, Sung P (2003). DNA helicase Srs2 disrupts the Rad51 presynaptic filament. **Nature** 423(6937): 305-309. doi: 10.1038/nature01577
89. Kaniecki K, De Tullio L, Gibb B, Kwon Y, Sung P, Greene EC (2017). Dissociation of Rad51 presynaptic complexes and heteroduplex DNA joints by tandem assemblies of Srs2. **Cell Rep** 21(11): 3166-3177. doi: 10.1016/j.celrep.2017.11.047
90. Fung CW, Mozlin AM, Symington LS (2009). Suppression of the double-strand-break-repair defect of the *Saccharomyces cerevisiae* rad57 mutant. **Genetics** 181(4): 1195-1206. doi: 10.1534/genetics.109.100842
91. St Onge RP, Mani R, Oh J, Proctor M, Fung E, Davis RW, Nislow C, Roth FP, Giaever G (2007). Systematic pathway analysis using high-resolution fitness profiling of combinatorial gene deletions. **Nature genetics** 39(2): 199-206. doi: 10.1038/ng1948
92. Chun J, Buechelmaier ES, Powell SN (2013). Rad51 paralogs BCDX2 and CX3 act at different stages in the BRCA1-BRCA2-dependent homologous recombination pathway. **Mol Cell Biol** 33(2): 387-395. doi: 10.1128/MCB.00465-12
93. Thacker J (1999). A surfeit of RAD51-like genes? **Trends Genet** 15(5): 166-168. doi: 10.1016/s0168-9525(99)01733-3
94. Braybrooke JP, Spink KG, Thacker J, Hickson ID (2000). The RAD51 family member, RAD51L3, is a DNA-stimulated ATPase that forms a complex with XRCC2. **J Biol Chem** 275(37): 29100-29106. doi: 10.1074/jbc.M002075200
95. Tsutsui Y, Morishita T, Iwasaki H, Toh H, Shinagawa H (2000). A recombination repair gene of *Schizosaccharomyces pombe*, rhp57, is a functional homolog of the *Saccharomyces cerevisiae* RAD57 gene and is phylogenetically related to the human XRCC3 gene. **Genetics** 154(4): 1451-1461. PMID: 10747044
96. Masson JY, Tarsounas MC, Stasiak AZ, Stasiak A, Shah R, McIlwraith MJ, Benson FE, West SC (2001). Identification and purification of two distinct complexes containing the five RAD51 paralogs. **Genes Dev** 15(24): 3296-3307. doi: 10.1101/gad.947001
97. Rodrigue A, Lafrance M, Gauthier MC, McDonald D, Hendzel M, West SC, Jasin M, Masson JY (2006). Interplay between human DNA repair proteins at a unique double-strand break in vivo. **EMBO J** 25(1): 222-231. doi: 10.1038/sj.emboj.7600914
98. Liu Y, Tarsounas M, O'Regan P, West SC (2007). Role of RAD51C and XRCC3 in genetic recombination and DNA repair. **J Biol Chem** 282(3): 1973-1979. doi: 10.1074/jbc.M609066200
99. Shor E, Weinstein J, Rothstein R (2005). A genetic screen for top3 suppressors in *Saccharomyces cerevisiae* identifies SHU1, SHU2, PSY3 and CSM2: four genes involved in error-free DNA repair. **Genetics** 169(3): 1275-1289. doi: 10.1534/genetics.104.036764
100. Sasanuma H, Tawaramoto MS, Lao JP, Hosaka H, Sanda E, Suzuki M, Yamashita E, Hunter N, Shinohara M, Nakagawa A, Shinohara A (2013). A new protein complex promoting the assembly of Rad51 filaments. **Nat Commun** 4:1676. doi: 10.1038/ncomms2678
101. Gaines WA, Godin SK, Kabbinar FF, Rao T, VanDemark AP, Sung P, Bernstein KA (2015). Promotion of presynaptic filament assembly by the ensemble of *S. cerevisiae* Rad51 paralogues with Rad52. **Nat Commun** 6:7834. doi: 10.1038/ncomms8834
102. Bernstein KA, Reid RJ, Sunjevaric I, Demuth K, Burgess RC, Rothstein R (2011). The Shu complex, which contains Rad51 paralogues, promotes DNA repair through inhibition of the Srs2 anti-recombinase. **Mol Biol Cell** 22(9): 1599-1607. doi: 10.1091/mbc.E10-08-0691
103. Uetz P, Giot L, Cagney G, Mansfield TA, Judson RS, Knight JR, Lockshon D, Narayan V, Srinivasan M, Pochart P, Qureshi-Emili A, Li Y, Godwin B, Conover D, Kalbfleisch T, Vijayadmodar G, Yang M, Johnston M, Fields S, Rothberg JM (2000). A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. **Nature** 403(6770): 623-627. doi: 10.1038/35001009
104. Liu T, Wan L, Wu Y, Chen J, Huang J (2011). hSWS1.SWSAP1 is an evolutionarily conserved complex required for efficient homologous recombination repair. **J Biol Chem** 286(48): 41758-41766. doi: 10.1074/jbc.M111.271080
105. Martin V, Chahwan C, Gao H, Blais V, Wohlschlegel J, Yates JR, 3rd, McGowan CH, Russell P (2006). Sws1 is a conserved regulator of homologous recombination in eukaryotic cells. **EMBO J** 25(11): 2564-2574. doi: 10.1038/sj.emboj.7601141
106. Mazin AV, Alexeev AA, Kowalczykowski SC (2003). A novel function of Rad54 protein. Stabilization of the Rad51 nucleoprotein filament. **J Biol Chem** 278(16): 14029-14036. doi: 10.1074/jbc.M212779200
107. Wolner B, Peterson CL (2005). ATP-dependent and ATP-independent roles for the Rad54 chromatin remodeling enzyme during recombinational repair of a DNA double strand break. **J Biol Chem** 280(11): 10855-10860. doi: 10.1074/jbc.M414388200
108. Agarwal S, van Cappellen WA, Guenole A, Eppink B, Linsen SE, Meijering E, Houtsmuller A, Kanaar R, Essers J (2011). ATP-dependent and independent functions of Rad54 in genome maintenance. **J Cell Biol** 192(5): 735-750. doi: 10.1083/jcb.201011025
109. Solinger JA, Kiianitsa K, Heyer WD (2002). Rad54, a Swi2/Snf2-like recombinational repair protein, disassembles Rad51:dsDNA filaments. **Mol Cell** 10(5): 1175-1188. doi: 10.1016/s1097-2765(02)00743-8
110. Mason JM, Dusad K, Wright WD, Grubb J, Budke B, Heyer WD, Connell PP, Weichselbaum RR, Bishop DK (2015). RAD54 family translocases counter genotoxic effects of RAD51 in human tumor cells. **Nucleic Acids Res** 43(6): 3180-3196. doi: 10.1093/nar/gkv175
111. Spies J, Waizenegger A, Barton O, Surder M, Wright WD, Heyer WD, Lobrich M (2016). Nek1 Regulates Rad54 to Orchestrate Homologous Recombination and Replication Fork Stability. **Mol Cell** 62(6): 903-917. doi: 10.1016/j.molcel.2016.04.032
112. Hayase A, Takagi M, Miyazaki T, Oshiumi H, Shinohara M, Shinohara A (2004). A protein complex containing Mei5 and Sae3 promotes the assembly of the meiosis-specific RecA homolog Dmc1. **Cell** 119(7): 927-940. doi: 10.1016/j.cell.2004.10.031
113. Khasanov FK, Salakhova AF, Khasanova OS, Grishchuk AL, Chepurnaja OV, Korolev VG, Kohli J, Bashkirov VI (2008). Genetic analysis reveals different roles of *Schizosaccharomyces pombe* sfr1/dds20 in meiotic and mitotic DNA recombination and repair. **Curr Genet** 54(4): 197-211. doi: 10.1007/s00294-008-0212-z
114. Yuan J, Chen J (2011). The role of the human SWI5-MEI5 complex in homologous recombination repair. **J Biol Chem** 286(11): 9888-9893. doi: 10.1074/jbc.M110.207290
115. Tsai SP, Su GC, Lin SW, Chung CI, Xue X, Dunlop MH, Akamatsu Y, Jasin M, Sung P, Chi P (2012). Rad51 presynaptic filament stabilization function of the mouse Swi5-Sfr1 heterodimeric complex. **Nucleic Acids Res** 40(14): 6558-6569. doi: 10.1093/nar/gks305
116. Kokabu Y, Murayama Y, Kuwabara N, Oroguchi T, Hashimoto H, Tsutsui Y, Nozaki N, Akashi S, Unzai S, Shimizu T, Iwasaki H, Sato M, Ikeguchi M (2011). Fission yeast Swi5-Sfr1 protein complex, an activator of Rad51 recombinase, forms an extremely elongated dogleg-shaped structure. **J Biol Chem** 286(50): 43569-43576. doi: 10.1074/jbc.M111.303339
117. Haruta N, Kurokawa Y, Murayama Y, Akamatsu Y, Unzai S, Tsutsui Y, Iwasaki H (2006). The Swi5-Sfr1 complex stimulates Rhp51/Rad51-

- and Dmc1-mediated DNA strand exchange in vitro. **Nat Struct Mol Biol** 13(9): 823-830. doi: 10.1038/nsmb11136
118. Alatwi HE, Downs JA (2015). Removal of H2A.Z by INO80 promotes homologous recombination. **EMBO Rep** 16(8): 986-994. doi: 10.15252/embr.201540330
119. Lademann CA, Renkawitz J, Pfander B, Jentsch S (2017). The INO80 complex removes H2A.Z to promote presynaptic filament formation during homologous recombination. **Cell Rep** 19(7): 1294-1303. doi: 10.1016/j.celrep.2017.04.051
120. Tsukuda T, Fleming AB, Nickloff JA, Osley MA (2005). Chromatin remodelling at a DNA double-strand break site in *Saccharomyces cerevisiae*. **Nature** 438(7066): 379-383. doi: 10.1038/nature04148
121. Chen L, Cai Y, Jin J, Florens L, Swanson SK, Washburn MP, Conaway JW, Conaway RC (2011). Subunit organization of the human INO80 chromatin remodeling complex: an evolutionarily conserved core complex catalyzes ATP-dependent nucleosome remodeling. **J Biol Chem** 286(13): 11283-11289. doi: 10.1074/jbc.M111.222505
122. Papamichos-Chronakis M, Krebs JE, Peterson CL (2006). Interplay between Ino80 and Swr1 chromatin remodeling enzymes regulates cell cycle checkpoint adaptation in response to DNA damage. **Genes Dev** 20(17): 2437-2449. doi: 10.1101/gad.1440206
123. van Attikum H, Fritsch O, Gasser SM (2007). Distinct roles for SWR1 and INO80 chromatin remodeling complexes at chromosomal double-strand breaks. **EMBO J** 26(18): 4113-4125. doi: 10.1038/sj.emboj.7601835
124. Shah PP, Zheng X, Epshtein A, Carey JN, Bishop DK, Klein HL (2010). Swi2/Snf2-related translocases prevent accumulation of toxic Rad51 complexes during mitotic growth. **Mol Cell** 39(6): 862-872. doi: 10.1016/j.molcel.2010.08.028
125. Chi P, Kwon Y, Seong C, Epshtein A, Lam I, Sung P, Klein HL (2006). Yeast recombination factor Rdh54 functionally interacts with the Rad51 recombinase and catalyzes Rad51 removal from DNA. **J Biol Chem** 281(36): 26268-26279. doi: 10.1074/jbc.M602983200
126. Raschle M, Van Komen S, Chi P, Ellenberger T, Sung P (2004). Multiple interactions with the Rad51 recombinase govern the homologous recombination function of Rad54. **J Biol Chem** 279(50): 51973-51980. doi: 10.1074/jbc.M410101200
127. Santa Maria SR, Kwon Y, Sung P, Klein HL (2013). Characterization of the interaction between the *Saccharomyces cerevisiae* Rad51 recombinase and the DNA translocase Rdh54. **J Biol Chem** 288(30): 21999-22005. doi: 10.1074/jbc.M113.480475
128. Chi P, Kwon Y, Visnapuu ML, Lam I, Santa Maria SR, Zheng X, Epshtein A, Greene EC, Sung P, Klein HL (2011). Analyses of the yeast Rad51 recombinase A265V mutant reveal different in vivo roles of Swi2-like factors. **Nucleic Acids Res** 39(15): 6511-6522. doi: 10.1093/nar/gkr297
129. Mazin AV, Mazina OM, Bugreev DV, Rossi MJ (2010). Rad54, the motor of homologous recombination. **DNA Repair (Amst)** 9(3): 286-302. doi: 10.1016/j.dnarep.2009.12.006
130. Boehm EM, Gildenberg MS, Washington MT (2016). The Many Roles of PCNA in Eukaryotic DNA Replication. **The Enzymes** 39:231-254. doi: 10.1016/bs.enz.2016.03.003
131. Bochman ML, Sabouri N, Zakian VA (2010). Unwinding the functions of the Pif1 family helicases. **DNA Repair (Amst)** 9(3): 237-249. doi: 10.1016/j.dnarep.2010.01.008
132. Pfander B, Moldovan GL, Sacher M, Hoegge C, Jentsch S (2005). SUMO-modified PCNA recruits Srs2 to prevent recombination during S phase. **Nature** 436(7049): 428-433. doi: 10.1038/nature03665
133. Burgess RC, Lisby M, Altmannova V, Krejci L, Sung P, Rothstein R (2009). Localization of recombination proteins and Srs2 reveals anti-recombinase function in vivo. **J Cell Biol** 185(6): 969-981. doi: 10.1083/jcb.200810055
134. Armstrong AA, Mohideen F, Lima CD (2012). Recognition of SUMO-modified PCNA requires tandem receptor motifs in Srs2. **Nature** 483(7387): 59-63. doi: 10.1038/nature10883
135. Aguilera A, Klein HL (1988). Genetic control of intrachromosomal recombination in *Saccharomyces cerevisiae*. I. Isolation and genetic characterization of hyper-recombination mutations. **Genetics** 119(4): 779-790. PMID: 3044923
136. Colavito S, Macris-Kiss M, Seong C, Gleeson O, Greene EC, Klein HL, Krejci L, Sung P (2009). Functional significance of the Rad51-Srs2 complex in Rad51 presynaptic filament disruption. **Nucleic Acids Res** 37(20): 6754-6764. doi: 10.1093/nar/gkp748
137. Branzei D, Vanoli F, Foiani M (2008). SUMOylation regulates Rad18-mediated template switch. **Nature** 456(7224): 915-920. doi: 10.1038/nature07587
138. Urulangodi M, Sebesta M, Menolfi D, Szakal B, Sollier J, Sisakova A, Krejci L, Branzei D (2015). Local regulation of the Srs2 helicase by the SUMO-like domain protein Esc2 promotes recombination at sites of stalled replication. **Genes Dev** 29(19): 2067-2080. doi: 10.1101/gad.265629.115
139. Feng W, Jasin M (2017). BRCA2 suppresses replication stress-induced mitotic and G1 abnormalities through homologous recombination. **Nat Commun** 8(1): 525. doi: 10.1038/s41467-017-00634-0
140. Zellweger R, Dalcher D, Mutreja K, Berti M, Schmid JA, Herrador R, Vindigni A, Lopes M (2015). Rad51-mediated replication fork reversal is a global response to genotoxic treatments in human cells. **J Cell Biol** 208(5): 563-579. doi: 10.1083/jcb.201406099
141. Hashimoto Y, Ray Chaudhuri A, Lopes M, Costanzo V (2010). Rad51 protects nascent DNA from Mre11-dependent degradation and promotes continuous DNA synthesis. **Nat Struct Mol Biol** 17(11): 1305-1311. doi: 10.1038/nsmb.1927
142. Kolinjivadi AM, Sannino V, De Antoni A, Zadorozhny K, Kilkenny M, Techer H, Baldi G, Shen R, Ciccio A, Pellegrini L, Krejci L, Costanzo V (2017). Smarcal1-Mediated Fork Reversal Triggers Mre11-Dependent Degradation of Nascent DNA in the Absence of Brca2 and Stable Rad51 Nucleofilaments. **Mol Cell** 67(5): 867-881 e867. doi: 10.1016/j.molcel.2017.07.001
143. Schlacher K, Christ N, Siau N, Egashira A, Wu H, Jasin M (2011). Double-strand break repair-independent role for BRCA2 in blocking stalled replication fork degradation by MRE11. **Cell** 145(4): 529-542. doi: 10.1016/j.cell.2011.03.041
144. Dungrawala H, Bhat KP, Le Meur R, Chazin WJ, Ding X, Sharan SK, Wessel SR, Sathe AA, Zhao R, Cortez D (2017). RADX Promotes Genome Stability and Modulates Chemosensitivity by Regulating RAD51 at Replication Forks. **Mol Cell** 67(3): 374-386 e375. doi: 10.1016/j.molcel.2017.06.023
145. Gonzalez-Prieto R, Munoz-Cabello AM, Cabello-Lobato MJ, Prado F (2013). Rad51 replication fork recruitment is required for DNA damage tolerance. **EMBO J** 32(9): 1307-1321. doi: 10.1038/emboj.2013.73
146. Tsurimoto T, Stillman B (1990). Functions of replication factor C and proliferating-cell nuclear antigen: functional similarity of DNA polymerase accessory proteins from human cells and bacteriophage T4. **Proc Natl Acad Sci USA** 87(3): 1023-1027. doi: 10.1073/pnas.87.3.1023
147. Cai J, Uhlmann F, Gibbs E, Flores-Rozas H, Lee CG, Phillips B, Finkelstein J, Yao N, O'Donnell M, Hurwitz J (1996). Reconstitution of human replication factor C from its five subunits in baculovirus-

- infected insect cells. **Proc Natl Acad Sci USA** 93(23): 12896-12901. doi: 10.1073/pnas.93.23.12896
148. Hingorani MM, Coman MM (2002). On the specificity of interaction between the *Saccharomyces cerevisiae* clamp loader replication factor C and primed DNA templates during DNA replication. **J Biol Chem** 277(49): 47213-47224. doi: 10.1074/jbc.M206764200
149. Tsurimoto T, Stillman B (1991). Replication factors required for SV40 DNA replication in vitro. II. Switching of DNA polymerase alpha and delta during initiation of leading and lagging strand synthesis. **J Biol Chem** 266(3): 1961-1968. PMID: 1671046
150. Gomes XV, Burgers PM (2001). ATP utilization by yeast replication factor C. I. ATP-mediated interaction with DNA and with proliferating cell nuclear antigen. **J Biol Chem** 276(37): 34768-34775. doi: 10.1074/jbc.M011631200
151. Waga S, Stillman B (1998). Cyclin-dependent kinase inhibitor p21 modulates the DNA primer-template recognition complex. **Mol Cell Biol** 18(7): 4177-4187. doi: 10.1128/mcb.18.7.4177
152. Li J, Holzschu DL, Sugiyama T (2013). PCNA is efficiently loaded on the DNA recombination intermediate to modulate polymerase delta, eta, and zeta activities. **Proc Natl Acad Sci U S A** 110(19): 7672-7677. doi: 10.1073/pnas.1222241110
153. Yuzhakov A, Kelman Z, Hurwitz J, O'Donnell M (1999). Multiple competition reactions for RPA order the assembly of the DNA polymerase delta holoenzyme. **EMBO J** 18(21): 6189-6199. doi: 10.1093/emboj/18.21.6189
154. De Tullio L, Kaniecki K, Kwon Y, Crickard JB, Sung P, Greene EC (2017). Yeast Srs2 Helicase Promotes Redistribution of Single-Stranded DNA-Bound RPA and Rad52 in Homologous Recombination Regulation. **Cell Rep** 21(3): 570-577. doi: 10.1016/j.celrep.2017.09.073
155. Ira G, Malkova A, Liberi G, Foiani M, Haber JE (2003). Srs2 and Sgs1-Top3 suppress crossovers during double-strand break repair in yeast. **Cell** 115(4): 401-411. doi: 10.1016/s0092-8674(03)00886-9
156. Miura T, Shibata T, Kusano K (2013). Putative antirecombinase Srs2 DNA helicase promotes noncrossover homologous recombination avoiding loss of heterozygosity. **Proc Natl Acad Sci USA** 110(40): 16067-16072. doi: 10.1073/pnas.1303111110
157. Liu J, Ede C, Wright WD, Gore SK, Jenkins SS, Freudenthal BD, Todd Washington M, Veaute X, Heyer WD (2017). Srs2 promotes synthesis-dependent strand annealing by disrupting DNA polymerase delta-extending D-loops. **Elife** 6: e22195. doi: 10.7554/eLife.22195
158. Yeung M, Durocher D (2011). Srs2 enables checkpoint recovery by promoting disassembly of DNA damage foci from chromatin. **DNA Repair (Amst)** 10(12): 1213-1222. doi: 10.1016/j.dnarep.2011.09.005
159. Vaze MB, Pelliccioli A, Lee SE, Ira G, Liberi G, Arbel-Eden A, Foiani M, Haber JE (2002). Recovery from checkpoint-mediated arrest after repair of a double-strand break requires Srs2 helicase. **Mol Cell** 10(2): 373-385. doi: 10.1016/s1097-2765(02)00593-2
160. van der Heijden T, Seidel R, Modesti M, Kanaar R, Wyman C, Dekker C (2007). Real-time assembly and disassembly of human RAD51 filaments on individual DNA molecules. **Nucleic Acids Res** 35(17): 5646-5657. doi: 10.1093/nar/gkm629
161. Li X, Heyer WD (2009). RAD54 controls access to the invading 3'-OH end after RAD51-mediated DNA strand invasion in homologous recombination in *Saccharomyces cerevisiae*. **Nucleic Acids Res** 37(2): 638-646. doi: 10.1093/nar/gkn980
162. Inano S, Sato K, Katsuki Y, Kobayashi W, Tanaka H, Nakajima K, Nakada S, Miyoshi H, Knies K, Takaori-Kondo A, Schindler D, Ishiai M, Kurumizaka H, Takata M (2017). RFD3-Mediated Ubiquitination Promotes Timely Removal of Both RPA and RAD51 from DNA Damage Sites to Facilitate Homologous Recombination. **Mol Cell** 66(5): 622-634 e628. doi: 10.1016/j.molcel.2017.04.022
163. Bugreev DV, Yu X, Egelman EH, Mazin AV (2007). Novel pro- and anti-recombination activities of the Bloom's syndrome helicase. **Genes Dev** 21(23): 3085-3094. doi: 10.1101/gad.1609007
164. Hu Y, Raynard S, Sehorn MG, Lu X, Bussen W, Zheng L, Stark JM, Barnes EL, Chi P, Janscak P, Jasin M, Vogel H, Sung P, Luo G (2007). RECQL5/Recq15 helicase regulates homologous recombination and suppresses tumor formation via disruption of Rad51 presynaptic filaments. **Genes Dev** 21(23): 3073-3084. doi: 10.1101/gad.1609107
165. Simandlova J, Zagelbaum J, Payne MJ, Chu WK, Shevelev I, Hanada K, Chatterjee S, Reid DA, Liu Y, Janscak P, Rothenberg E, Hickson ID (2013). FBH1 helicase disrupts RAD51 filaments in vitro and modulates homologous recombination in mammalian cells. **J Biol Chem** 288(47): 34168-34180. doi: 10.1074/jbc.M113.484493
166. Sommers JA, Rawtani N, Gupta R, Bugreev DV, Mazin AV, Cantor SB, Brosh RM, Jr. (2009). FANCD1 uses its motor ATPase to destabilize protein-DNA complexes, unwind triplexes, and inhibit RAD51 strand exchange. **J Biol Chem** 284(12): 7505-7517. doi: 10.1074/jbc.M809019200
167. Moldovan GL, Dejsuphong D, Petalcorin MI, Hofmann K, Takeda S, Boulton SJ, D'Andrea AD (2012). Inhibition of homologous recombination by the PCNA-interacting protein PARI. **Mol Cell** 45(1): 75-86. doi: 10.1016/j.molcel.2011.11.010
168. Fugger K, Mistrik M, Danielsen JR, Dinant C, Falck J, Bartek J, Lukas J, Mailand N (2009). Human Fbh1 helicase contributes to genome maintenance via pro- and anti-recombinase activities. **J Cell Biol** 186(5): 655-663. doi: 10.1083/jcb.200812138
169. Traverso G, Bettegowda C, Kraus J, Speicher MR, Kinzler KW, Vogelstein B, Lengauer C (2003). Hyper-recombination and genetic instability in BLM-deficient epithelial cells. **Cancer Res** 63(24): 8578-8581. PMID: 14695165
170. Hu Y, Lu X, Barnes E, Yan M, Lou H, Luo G (2005). Recq15 and Blm RecQ DNA helicases have nonredundant roles in suppressing crossovers. **Mol Cell Biol** 25(9): 3431-3442. doi: 10.1128/MCB.25.9.3431-3442.2005
171. Lydeard JR, Lipkin-Moore Z, Jain S, Eapen VV, Haber JE (2010). Sgs1 and exo1 redundantly inhibit break-induced replication and de novo telomere addition at broken chromosome ends. **PLoS genetics** 6(5): e1000973. doi: 10.1371/journal.pgen.1000973
172. Chung WH, Zhu Z, Papusha A, Malkova A, Ira G (2010). Defective resection at DNA double-strand breaks leads to de novo telomere formation and enhances gene targeting. **PLoS genetics** 6(5): e1000948. doi: 10.1371/journal.pgen.1000948
173. Barlow JH, Rothstein R (2009). Rad52 recruitment is DNA replication independent and regulated by Cdc28 and the Mec1 kinase. **EMBO J** 28(8): 1121-1130. doi: 10.1038/emboj.2009.43
174. Lisby M, Rothstein R, Mortensen UH (2001). Rad52 forms DNA repair and recombination centers during S phase. **Proc Natl Acad Sci U S A** 98(15): 8276-8282. doi: 10.1073/pnas.121006298
175. Holt LJ, Tuch BB, Villen J, Johnson AD, Gygi SP, Morgan DO (2009). Global analysis of Cdk1 substrate phosphorylation sites provides insights into evolution. **Science** 325(5948): 1682-1686. doi: 10.1126/science.1172867
176. Sacher M, Pfander B, Hoegge C, Jentsch S (2006). Control of Rad52 recombination activity by double-strand break-induced SUMO modification. **Nature cell biology** 8(11): 1284-1290. doi: 10.1038/ncb1488
177. Altmanova V, Eckert-Boulet N, Arneric M, Kolesar P, Chaloupkova R, Damborsky J, Sung P, Zhao X, Lisby M, Krejci L (2010).

Rad52 SUMOylation affects the efficiency of the DNA repair. **Nucleic Acids Res** 38(14): 4708-4721. doi: 10.1093/nar/gkq195

178. Bashkirov VI, King JS, Bashkirova EV, Schmuckli-Maurer J, Heyer WD (2000). DNA repair protein Rad55 is a terminal substrate of the DNA damage checkpoints. **Mol Cell Biol** 20(12): 4393-4404. doi: 10.1128/mcb.20.12.4393-4404.2000

179. Herzberg K, Bashkirov VI, Rolfsmeier M, Haghazari E, McDonald WH, Anderson S, Bashkirova EV, Yates JR, 3rd, Heyer WD (2006). Phosphorylation of Rad55 on serines 2, 8, and 14 is required for efficient homologous recombination in the recovery of stalled replication forks. **Mol Cell Biol** 26(22): 8396-8409. doi: 10.1128/MCB.01317-06

180. Flott S, Kwon Y, Pigli YZ, Rice PA, Sung P, Jackson SP (2011). Regulation of Rad51 function by phosphorylation. **EMBO Rep** 12(8): 833-839. doi: 10.1038/embor.2011.127

181. Dou H, Huang C, Singh M, Carpenter PB, Yeh ET (2010). Regulation of DNA repair through deSUMOylation and SUMOylation of replication protein A complex. **Mol Cell** 39(3): 333-345. doi: 10.1016/j.molcel.2010.07.021

182. Cremona CA, Sarangi P, Yang Y, Hang LE, Rahman S, Zhao X (2012). Extensive DNA damage-induced sumoylation contributes to replication and repair and acts in addition to the mec1 checkpoint. **Mol Cell** 45(3): 422-432. doi: 10.1016/j.molcel.2011.11.028

183. Psakhye I, Jentsch S (2012). Protein group modification and synergy in the SUMO pathway as exemplified in DNA repair. **Cell** 151(4): 807-820. doi: 10.1016/j.cell.2012.10.021

184. Shima H, Suzuki H, Sun J, Kono K, Shi L, Kinomura A, Horikoshi Y, Ikura T, Ikura M, Kanaar R, Igarashi K, Saitoh H, Kurumizaka H, Tashiro S (2013). Activation of the SUMO modification system is required for the accumulation of RAD51 at sites of DNA damage. **J Cell Sci** 126(Pt 22): 5284-5292. doi: 10.1242/jcs.133744

185. Luo K, Li L, Li Y, Wu C, Yin Y, Chen Y, Deng M, Nowsheen S, Yuan J, Lou Z (2016). A phosphorylation-deubiquitination cascade regulates the BRCA2-RAD51 axis in homologous recombination. **Genes Dev** 30(23): 2581-2595. doi: 10.1101/gad.289439.116

186. Liu J, Kruswick A, Dang H, Tran AD, Kwon SM, Wang XW, Oberdoerffer P (2017). Ubiquitin-specific protease 21 stabilizes BRCA2 to control DNA repair and tumor growth. **Nat Commun** 8(1): 137. doi: 10.1038/s41467-017-00206-2

187. Chiolo I, Saponaro M, Baryshnikova A, Kim JH, Seo YS, Liberi G (2007). The human F-box DNA helicase FBH1 faces *Saccharomyces cerevisiae* Srs2 and postreplication repair pathway roles. **Mol Cell Biol** 27(21): 7439-7450. doi: 10.1128/MCB.00963-07

188. Saponaro M, Callahan D, Zheng X, Krejci L, Haber JE, Klein HL, Liberi G (2010). Cdk1 targets Srs2 to complete synthesis-dependent strand annealing and to promote recombinational repair. **PLoS genetics** 6(2): e1000858. doi: 10.1371/journal.pgen.1000858

189. Kolesar P, Sarangi P, Altmannova V, Zhao X, Krejci L (2012). Dual roles of the SUMO-interacting motif in the regulation of Srs2 sumoylation. **Nucleic Acids Res** 40(16): 7831-7843. doi: 10.1093/nar/gks484